

## Acceptable Collection Container

## **Urine Drug Screen**

| Test Includes:                     |                   | Amphetamine Barbiturates Benzodiazepines Buprenorphine Cocaine Methamphetamine Methadone Opiates Oxycodone Phencyclidine Propoxyphene Cannabinoids Tricyclic Antidepressants   |
|------------------------------------|-------------------|--|
| Performing Lab:                    |                   | New Hanover and Cape Fear  |
| Specimen Container:                |                   | Clean screw-cap container  |
| Minimum Volume Required:           |                   | 1.0 mL   |
| Testing Availability               | Routine:<br>Stat: | 24 hours/day<br>N/A  |
| Turnaround Time:                   |                   | Routine: 4 hours   |
| Special Handling:                  |                   | None   |
| Patient Preparation:               |                   | None   |
| Specimen Stability:                |                   | 48 hours at 2-8 degrees C.   |
| Reference Range:                   |                   | None   |
| Cutoff concentration for positive: |                   | Amphetamine: 500 ng/mL Barbiturates: 200 ng/mL Benzodiazepines: 150 ng/mL Buprenorphine: 10 ng/mL Cocaine: 150 ng/mL Methamphetamine: 500 ng/mL Methadone: 200 ng/mL Opiates: 100 ng/mL Oxycodone: 100 ng/mL Phencyclidine: 25 ng/mL |

|                                | Propoxyphene: 300 ng/mL<br>Cannabinoids: 50 ng/mL<br>Tricyclic Antidepressants: 300 ng/mL |
|--------------------------------|---|
| CPT Code:                      | G0434   |
| Testing Methodology:           | One-step immunochromatographic  |
| Causes for Specimen Rejection: | Improper labeling   |
| Other Comments:                |   |
| Clinical Significance:         | For the qualitative detection of multiple drugs and drug metabolites in urine.            |